

References

- Barbas KH, Kelleher DK. Breastfeeding success among infants with congenital heart disease. *Pediatric Nursing*. 2004;30:285-89.
- Bartick M, Reinhold A. The burden of suboptimal breastfeeding in the United States: a pediatric cost analysis. *Pediatrics*. 2010;125:e1048–56.
- Bartick M, Steube A, Schwarz EB, et.al. Cost Analysis of Maternal Disease Associated With Suboptimal Breastfeeding. *Obstet Gynecol*. 2013;122(1):111–9.
- Bisquera JA, Cooper TR, Berseth CL. Impact of necrotizing enterocolitis on length of stay and hospital charges in very low birth weight infants. *Pediatrics*. 2002;109(3): 423-28.
- Caicedo R, Schanler R, Li N, Neu J. The developing eco-system: Implications for the neonate. *Pediatric Research* . 2005;58(4):625-28.
- Combs VL, Marino BL. A comparison of growth patterns in breast and bottle-fed infants with congenital heart disease. *Pediatric Nursing*. 1993;19:175-9.
- Cotterman, KJ. *Reverse Pressure Softening: A Simple Tool to Prepare Areola for Easier Latching During Engorgement*. *J Hum Lact*. 2004;20(2):227-37.
- Danner S, Cerutti E. Nursing your premature baby. [Childbirth Graphics](#) 1995:[Waco, Texas](#).
- Eidelman AI, Schanler RJ. **American Academy of Pediatrics, Section on Breastfeeding**. Breastfeeding and the use of human milk. *Pediatrics*. 2012; 129(3):2011-3552.
- Furman L, Taylor G, Minich N, Hack M. The effect of maternal milk on neonatal morbidity of very low-birth-weight infants. *Arch Pediatr Adolesc Med*. 2003 Jan; 157(1):66–71.
- Haase B, Barreira J, Murphy P, et. al. The Development of an Accurate Test Weighing Technique for Preterm and High Risk Hospitalized Infants. *BF Med*. 2009.

References

- Hurtekant, K. Spatz, D. Special Considerations for Breastfeeding the Infant with Spina Bifida. *J Perinat Neonat Nurs.* 2007; 21(1): 69-75.
- Ip S, Chung M, Raman G, et al. Breastfeeding and maternal and infant health outcomes in developed countries. In: *Evidence Report/ Technology Assessment 2007*;153:1-186. Rockville (MD). Agency for Healthcare Research and Quality (AHRQ), US Department of Health and Human Services.
- Marino BL, O'Brien P, LoRe H. Oxygen saturations during breast and bottle feedings in infants with congenital heart disease. *J Pediatr Nur.* 1995;10:360-4.
- Meier PP. Bottle- and breast-feeding: Effects on transcutaneous oxygen pressure and temperature in preterm infants. *Nurs Res.* 1988;37:36-41.
- Meier PP, Brown L, Hurst N, et al. Nipple shields for preterm infants: Effect on milk transfer and duration of breastfeeding. *J Hum Lact.* 2000;16:106-114.
- Meinzen-Derr J, Poindexter B, Wrage L, Morrow AL, Stoll B, Donovan EF. Role of human milk in extremely low birth weight infants' risk of necrotizing enterocolitis or death. *J Perinatol.* 2009;29(1):57–62.
- Miracle DJ, Meier PP, Bennett PA. Mother's decision to change from formula to mothers' milk for their very low birthweight infants. *JOGNN.* 2004;33:692-703.
- Morton J, Hall JY, Wong RJ, et al. Combining hand techniques with electric pumping increases milk production in mother's of preterm infants. *Journal of Perinatology.* 2009; 29:757–64.
- Patel AL, Meier PP, Engstrom JL. The evidence for Use of Human Milk in very low- birth weight preterm infants. *NeoReviews.* 2007; 8: e459-e466.

References

- Payne NR, Carpenter JH, Badger GJ, Horbar JD, Rogowski J. Marginal increase in cost and excess length of stay associated with nosocomial bloodstream infections in surviving very low birth weight infants. *Pediatrics*. 2004;114(2):348-55.
- Rodriguez N, Miracle DJ, Meier PP. Sharing the science on human milk feedings with mothers of very-low-birth-weight infants. *JOGNN*. 2005;34:109-119.
- Schanler RJ, Shulman RJ, Lau C. et al. Feeding Strategies for premature infants: beneficial outcomes of feeding fortified human milk vs. preterm formula. *Pediatrics*. 1999;103: 434-9.
- Schwarz EB, Brown JS, Creasman JM, et al. Lactation and Maternal Risk of Type 2 Diabetes: A Population-based Study. *The American Journal of Medicine*. 2010;123(9):863.e1-6.
- Schwarz EB, Ray RM, Stuebe AM, et al. Duration of lactation and risk factors for maternal cardiovascular disease. *Obstetrics and gynecology*. 2009;113(5):974-82.
- Sisk PM, Lovelady CA, Dillard RG, Gruber KJ, O'Shea TM. Early human milk feeding is associated with a lower risk of enterocolitis in very low birth weight infants. *J Perinatol*. 2007;27(7):428-33.
- Spatz D. State of the Science: Use of Human Milk and Breastfeeding for Vulnerable Infants. *J Perinat Neonat Nurs*. 2006; 20(1):51-5.
- Spatz D, Goldschmidt K. Preserving Breastfeeding for the Re-hospitalized Infant: A Clinical Pathway. *MCN*. 2006; 31(1):45-51.
- Sullivan S, Schanler RJ, Kim JH, Patel AL, Trawoger R, et al. An exclusively human milk-based diet is associated with a lower rate of necrotizing enterocolitis than a diet of human milk and bovine milk-based products. *J Pediatr*. 2010;156(4):562-7.